

This Page Is Inserted by IFW Operations
and is not a part of the Official Record

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images may include (but are not limited to):


- BLACK BORDERS
- TEXT CUT OFF AT TOP, BOTTOM OR SIDES
- FADED TEXT
- ILLEGIBLE TEXT
- SKEWED/SLANTED IMAGES
- COLORED PHOTOS
- BLACK OR VERY BLACK AND WHITE DARK PHOTOS
- GRAY SCALE DOCUMENTS

IMAGES ARE BEST AVAILABLE COPY.

**As rescanning documents *will not* correct images,
please do not report the images to the
Image Problem Mailbox.**


Claims

1. A glucose-solasodine conjugate of the general formula I or a derivative thereof



wherein each of R_1 and R_2 are the same or different and represents, R_1 is a benzoyl or a pivaloyl group.

2. A method for the preparation of the glucose-solasodine conjugate as defined in claim 1, comprising the reaction of solasodine with a glucoopyranosyl donor of general formula II




wherein each R_3 independently represents a benzoyl, acetyl or pivaloyl group,

wherein R_4 is halogen selected from Cl, Br or I and R_5 is hydrogen or


R_4 is hydrogen and R_5 is SEt or SPh,

followed by optionally de-protecting the obtained glycoside to yield a compound of the formula V

16




and reesterification of the most reactive hydroxyl groups (OH-3 and OH-6) to yield a compound of the formula IIa




wherein R_2 is a group selected from pivaloyl or acetyl.

3. A method for the preparation of solamargine comprising the glycosylation of the diol of formula IIa



wherein R_2 is defined as in claim 1 with an α -L-rhamnopyranosyl donor

to yield protected solamargine of formula III (1) which is de-esterified to yield solamargine of formula III (2)




(1) $R_1 = \text{Piv}$ and $R_2 = \text{Benzoyl or Acetyl}$

(2) $R_1 = R_2 = \text{H}$

4. The method according to claim 2, wherein the D-glucosepyranosyl donor is tetra-O-benzoyl- α -D-glucopyranosyl bromide, tetra-O-acetyl- α -D-glucopyranosyl bromide or tetra-O-pivaloyl- α -D-glucopyranosyl bromide.
5. The method according to claim 2 or 4, wherein the glycosylation reaction is carried out in the presence of a promoter selected from silver trifluoromethane sulfonate (silver triflate), boron trifluoride diethyl etherate, trimethylsilyl triflate bromide, N-iodosuccinimide or dimethyl thiomethyl sulfonium triflate, silver trifluoromethyltriflate.
6. The method of claim 2, wherein the protected glycoside is deprotected in methanol-dichloromethane solution by treatment with sodium methoxide, followed by neutralization with solid CO_2 or mild acid ion-exchange resin.
7. The method of claim 2, wherein the most reactive hydroxyl groups (OH-3 and OH-6) are protected by reesterification with pivaloyl chloride in pyridine solution.

8. The method of claim 3, wherein the rhamnose donor is tri-O-benzoyl- α -L-rhamnopyranosyl bromide, tri-O-pivaloyl- α -L-rhamnopyranosyl trichloroacetimidate or a glycoside of the general formula IV



IV

wherein R_6 is Br, Cl, I, SEt or SPh and

R_7 is benzoyl, acetyl or pivaloyl.

9. The method of claim 3, wherein the protected solamargine is de-esterified by treatment with a base selected from sodium methoxide or sodium hydroxide in methanol-dichloromethane solution or a methanol-tetrahydrofuran-water mixture followed by neutralization with solid CO_2 or mild acid ion-exchange resin.